



KWAZULU-NATAL PROVINCE

EDUCATION
REPUBLIC OF SOUTH AFRICA

**NATIONAL
SENIOR CERTIFICATE**

GRADE 11

MATHEMATICS PAPER 1

NOVEMBER 2023

MARKS: 150

TIME: 3 hours

Stanmorephysics

This question paper consists of 8 pages and 1 information sheet.

INSTRUCTIONS AND INFORMATION

Read the following instructions carefully before answering the questions.

- 
1. This question paper consists of 9 questions.
 2. Answer ALL the questions.
 3. Number the answers correctly according to the numbering system used in this question paper.
 4. Clearly show ALL calculations, diagrams, graphs, etc. that you have used in determining your answers.
 5. Answers only will NOT necessarily be awarded full marks.
 6. You may use an approved scientific calculator (non-programmable and non-graphical), unless stated otherwise.
 7. If necessary, round off answers to TWO decimal places, unless stated otherwise.
 8. Diagrams are NOT necessarily drawn to scale.
 9. An information sheet with formulae is included at the end of the question paper.
 10. Write neatly and legibly.

QUESTION 11.1 Solve for x :

$$1.1.1 \quad (2x+1)(5-x)=0 \quad (2)$$

$$1.1.2 \quad x^2 - \frac{2}{3}x - 5 = 0 \quad (\text{correct to TWO decimal places}) \quad (3)$$

$$1.1.3 \quad 2x\left(x - \frac{1}{2}\right) \leq 6 \quad (4)$$

$$1.1.4 \quad x + \sqrt{2x-3} = 3 \quad (5)$$

1.2 Solve for x and y simultaneously:

$$3x - y = 1 \quad \text{and} \quad 2xy - 3y^2 = -x^2 - 7 \quad (6)$$

1.3 The roots of a quadratic equation are $x = \frac{-10 \pm \sqrt{100 - 4k^2}}{2k}$.Calculate the values of k for which the roots are:1.3.1 equal (3)1.3.2 non-real (3)**[26]****QUESTION 2**

2.1 Simplify the following expressions completely, WITHOUT using a calculator:

$$2.1.1 \quad \frac{(9x^2)^{-\frac{1}{2}}}{9x^{-2}} \quad (3)$$

$$2.1.2 \quad \frac{\sqrt{8k^4} - \sqrt{98k^4}}{\sqrt{25k^2}} \quad (3)$$

2.2 Solve for x WITHOUT using a calculator:

$$2.2.1 \quad x^{\frac{3}{2}} = 125 \quad (3)$$

$$2.2.2 \quad \frac{p^x \cdot p^{\frac{2}{x}}}{p^{5-x}} = 1 \quad (4)$$

$$2.2.3 \quad \frac{x-4}{\sqrt{x-2}} = 5a^0 \quad (4)$$

[17]

QUESTION 3

Consider the following linear number pattern: $3x + 8$; $x - 2$; ... ; -141 ; -145 .

- 3.1 Is the above linear number pattern finite or infinite? (1)
- 3.2 Show that $x = -3$. (2)
- 3.3 Determine T_n , the general term of this number pattern, in the form $T_n = bn + c$. (3)
- 3.4 Calculate the value of T_{15} , the 15th term of this number pattern. (2)
- 3.5 How many terms are in this number pattern? (2)
- [10]**

QUESTION 4

Given a quadratic sequence: 3 ; 17 ; 35 ; 57 ; ...

- 4.1 Write down the next two terms of the sequence. (2)
- 4.2 Determine the general term of the sequence in the form $T_n = an^2 + bn + c$. (4)
- 4.3 Which term in the sequence is the second term greater than 826? (4)
- 4.4 Which TWO consecutive terms in the quadratic sequence have a first difference of 122? (4)
- 4.5 The given quadratic sequence is multiplied by a constant to obtain a new quadratic sequence: -6 ; $8p - 2$; -70 ; $18q$; ...
Write down the values of p and q . (2)
- [16]**

QUESTION 5

5.1 Given: $h(x) = \frac{-3}{x - \frac{3}{2}} - 2$



5.1.1 Write down the equations of asymptotes of h . (2)

5.1.2 Write down the coordinates of the y-intercept of h . (2)


5.1.3 Calculate the coordinates of the x-intercept of h . (3)

5.1.4 Draw the graph of h . Clearly show the asymptotes and the intercepts with the axes. (3)

5.1.5 Determine the equation of the axis of symmetry of h that has a negative gradient. (2)

5.1.6 Calculate the coordinates of the points of intersection of h and the axis of symmetry determined in QUESTION 5.1.5. (5)

5.2 The graph of $k(x) = b^x + q$ is drawn in the sketch below. The point $A(0 ; 1)$ lies on the asymptote of k and $B(1 ; 1,5)$ lies on k .



5.2.1 Determine the equation of k . (3)


5.2.2 Describe, in words, the transformation from k to $j(x) = 2^x$. (2)

[22]

QUESTION 6

The sketch below shows the graphs of $f(x) = \frac{1}{2}(x-2)^2 - 8$ and $g(x) = x - 6$.

- A and B are the x -intercepts of f .
- D is the turning point of f .
- C is the y -intercept of both graphs.
- Graphs f and g intersect at B and C.



- 6.1 Calculate the length of AB. (4)
- 6.2 Write down the coordinates of D. (1)
- 6.3 Write down the range of f . (1)
- 6.4 Calculate the average gradient between points A and C. (3)
- 6.5 For which values of k will $\frac{1}{2}x^2 - 2x - 6 - k = 0$ have two positive roots? (2)
- 6.6 Use the graphs to determine the values of x for which:
- 6.6.1 $f(x) \geq 0$ (2)
- 6.6.2 $\frac{g(x)}{f(x)} > 0$ (2)
- 6.7 Determine the maximum value of h if $h(x) = \sqrt{2^{-f(x)}}$ (3)
- 6.8 Calculate the value(s) of k for which $y = -3x + k$ will NOT be a tangent to f . (5)

[23]

QUESTION 7

- 7.1 A construction vehicle was bought for R625 000. The value of the vehicle depreciates annually at a rate of 10,25% p.a. on the reducing-balance method. Calculate the book value of the vehicle 6 years after it was bought. (2)
- 7.2 Calculate the effective interest rate if an investment earns interest at the rate of 8% p.a., compounded half-yearly. (3)
- 7.3 A house was bought $4\frac{1}{2}$ years ago. The value of the house increased at 14% p.a., compounded quarterly. If the value of the house is now R1 671 650,74, calculate the purchase price of the house. (3)
- 7.4 Mr Nsele invested R x into a savings account. He withdrew R15 000 from the account two years after making the initial deposit. Four years after making the initial deposit, the accumulated amount in the savings account was R75 000. The account earned interest at the rate of 12% p.a., compounded monthly for the first three years. Thereafter the interest rate changed to 8% p.a., compounded half-yearly.
- 7.4.1 Calculate the amount that Mr Nsele had in the account three years after making the initial deposit. (3)
- 7.4.2 Calculate the value of x , the amount Mr Nsele initially deposited into the account. (4)

[15]

QUESTION 8

8.1 For events A and B, it is given that $P(A) = 0,35$ and $P(B) = 0,4$.

8.1.1 Calculate $P(A \text{ or } B)$ if A and B are mutually exclusive. (2)

8.1.2 Calculate $P(A \text{ or } B)$ if A and B are independent. (3)



8.2 A survey was conducted among 150 Grade 11 learners about the genre of music that they listen to mostly. The results are shown in the table below.

	Classical	Pop	Hip hop	Total
Girls	35	k	27	80
Boys	32	15	23	70
Total	67		50	150

8.2.1 Write down the value of k . (1)

8.2.2 If a learner is selected at random from the group, what is the probability that the learner is a boy and does not listen to Pop music? (2)

8.2.3 Are the events: “being a girl” and “listening to Pop music” independent? Show all necessary calculations. (4)
[12]

QUESTION 9

9.1 The probability that South Africa (SA) will reach the finals in the 2027 Rugby World Cup is 0,35 and the probability that New Zealand (NZ) will reach the finals is 0,60. The probability that neither South Africa nor New Zealand will reach the final is 0,08.

Calculate the probability that only New Zealand will reach the finals. (4)

9.2 A picnic basket contains x apples and 8 bananas. A fruit is selected at random and eaten. Thereafter, another fruit is selected at random and eaten. The probability of selecting two different fruits, in any order, in the first two selections is $\frac{48}{95}$.

Calculate the total number of apples in the basket. (5)
[9]

TOTAL: 150

INFORMATION SHEET

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$A = P(1 + ni)$$

$$A = P(1 - ni)$$

$$A = P(1 - i)^n$$

$$A = P(1 + i)^n$$

$$T_n = a + (n - 1)d$$

$$S_n = \frac{n}{2} [2a + (n - 1)d]$$

$$T_n = ar^{n-1}$$

$$S_n = \frac{a(r^n - 1)}{r - 1}; r \neq 1$$

$$S_\infty = \frac{a}{1 - r}; -1 < r < 1$$

$$F = \frac{x[(1 + i)^n - 1]}{i}$$

$$P = \frac{x[1 - (1 + i)^{-n}]}{i}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h}$$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$$

$$y = mx + c$$

$$y - y_1 = m(x - x_1)$$

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

$$m = \tan \theta$$

$$(x - a)^2 + (y - b)^2 = r^2$$

$$\text{In } \triangle ABC: \quad \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\text{area } \triangle ABC = \frac{1}{2} ab \sin C$$

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

$$\cos 2\alpha = \begin{cases} \cos^2 \alpha - \sin^2 \alpha \\ 1 - 2\sin^2 \alpha \\ 2\cos^2 \alpha - 1 \end{cases}$$

$$\sin 2\alpha = 2\sin \alpha \cos \alpha$$

$$\bar{x} = \frac{\sum x}{n}$$

$$\sigma^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}$$

$$P(A) = \frac{n(A)}{n(S)}$$

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$

$$\hat{y} = a + bx$$

$$b = \frac{\sum (x - \bar{x})(y - \bar{y})}{\sum (x - \bar{x})^2}$$



KWAZULU-NATAL PROVINCE

EDUCATION
REPUBLIC OF SOUTH AFRICA

NATIONAL SENIOR CERTIFICATE

GRADE 11

MATHEMATICS PAPER 1

NOVEMBER 2023

MARKING GUIDELINES




MARKS: 150

Stanmorephysics



These marking guidelines consists of 17 pages.

CAPS-GRADE 11
Marking Guidelines

QUESTION 1

1.1.1	$(2x+1)(5-x)=0$ $x = -\frac{1}{2}$ or $x=5$ 	✓ answer ✓ answer (2)
1.1.2	$x^2 - \frac{2}{3}x - 5 = 0$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $x = \frac{-\left(-\frac{2}{3}\right) \pm \sqrt{\left(-\frac{2}{3}\right)^2 - 4(1)(-5)}}{2(1)}$ $x = 2,59$ or $x = -1,93$	✓ substitution into correct formula ✓ answer ✓ answer (3)
1.1.3	$2x\left(x - \frac{1}{2}\right) \leq 6$ $2x^2 - 1x - 6 \leq 0$ $(2x+3)(x-2) \leq 0$   $-\frac{3}{2} \leq x \leq 2$ OR $x \in \left[-1\frac{1}{2}; 2\right]$	✓ standard form ✓ factors ✓ ✓ $-\frac{3}{2} \leq x \leq 2$ (4)
1.1.4	$x + \sqrt{2x-3} = 3$ $\sqrt{2x-3} = 3-x$ $(\sqrt{2x-3})^2 = (3-x)^2$ $2x-3 = 9-6x+x^2$ $x^2 - 8x + 12 = 0$ $(x-6)(x-2) = 0$ $x \neq 6$ or $x = 2$	✓ isolating the root ✓ squaring both sides ✓ standard form ✓ $x \neq 6$ ✓ $x = 2$ (5)

CAPS-GRADE 11
Marking Guidelines

1.2	$3x - y = 1$ and $2xy - 3y^2 = -x^2 - 7$ $y = 3x - 1 \dots\dots\dots(3)$ $2x(3x - 1) - 3(3x - 1)^2 = -x^2 - 7$ $6x^2 - 2x - 27x^2 + 18x - 3 + x^2 + 7 = 0$ $-20x^2 + 16x + 4 = 0$ $5x^2 - 4x - 1 = 0$ $(5x + 1)(x - 1) = 0$ $x = \frac{-1}{5}$ or $x = 1$ $y = 3\left(\frac{-1}{5}\right) - 1$ $y = 3(1) - 1$ $y = \frac{-8}{5}$ or $y = 2$	equation 1 & equation 2 resp. ✓ equation 3 ✓ substitution into equation 2 ✓ standard form ✓ factors ✓ both x-values ✓ both y-values (6)
1.3	$x = \frac{-10 \pm \sqrt{100 - 4k^2}}{2k}$	
1.3.1	$100 - 4k^2 = 0$ $-4(k^2 - 25) = 0$ $(k + 5)(k - 5) = 0$ $k = 5$ or $k = -5$ OR $100 - 4k^2 = 0$ $4k^2 = 100$ $k^2 = 25$ $k = \pm 5$	✓ $100 - 4k^2 = 0$ ✓ factors ✓ both answers for k (3) OR ✓ $100 - 4k^2 = 0$ ✓ isolating the square ✓ both answers for k (3)
1.3.2	$100 - 4k^2 < 0$ $(k + 5)(k - 5) > 0$ CV: $k = -5$ or $k = 5$   $k < -5$ or $k > 5$ OR $k \in (-\infty; -5) \cup (5; \infty)$	✓ $100 - 4k^2 < 0$ ✓✓ $k < -5$ or $k > 5$ (3)
		[26]

CAPS GRADE 11
Marking Guidelines

QUESTION 2

2.1.1	$\frac{(9x^2)^{-\frac{1}{2}}}{9x^{-2}}$  $= \frac{3^{2\left(\frac{-1}{2}\right)} \cdot x^{2\left(\frac{-1}{2}\right)}}{3^2 \cdot x^{-2}}$ $= \frac{3^{-1} \cdot x^{-1}}{3^2 \cdot x^{-2}}$ $= 3^{-1-2} \cdot x^{-1+2}$ $= 3^{-3} \cdot x^1$ $= \frac{1}{27} x$ OR $\frac{(9x^2)^{-\frac{1}{2}}}{9x^{-2}}$ $= \frac{x^2}{9(9x^2)^{\frac{1}{2}}}$ $= \frac{x^2}{9(3x)}$ $\frac{1}{27} \text{ or } \frac{1}{27} x$	✓ splitting to prime bases & removing brackets around bases ✓ applying exponential law ✓ answer (3) OR ✓ changing to positive exponents ✓ applying exponential law ✓ answer (3)
2.1.2	$\sqrt{8k^4} - \sqrt{98k^4}$ $\sqrt{2^3} k^2$ $\sqrt{8} \cdot \sqrt{k^4} - \sqrt{98} \cdot \sqrt{k^4}$ $\sqrt{25k^2}$ $= \frac{\sqrt{4 \times 2 \cdot k^2} - \sqrt{49 \times 2 \cdot k^2}}{5k^2}$ $= \frac{2\sqrt{2}k^2 - 7\sqrt{2}k^2}{5k^2}$ $= \frac{-5\sqrt{2}k^2}{5k^2}$ $= -\sqrt{2}$	✓ applying rule: $\sqrt{ab} = \sqrt{a} \cdot \sqrt{b}$ to simplify the surds ✓ $\frac{-5\sqrt{2}k^2}{5k^2}$ ✓ answer (3)

CAPS-GRADE 11
Marking Guidelines

2.2.1	$x^{\frac{3}{2}} = 125$ $x^{\frac{3}{2}} = 5^3$ $x^{\frac{3}{2} \left(-\frac{2}{3} \right)} = 5^{3 \left(-\frac{2}{3} \right)}$ $x = 5^{-2}$ $x = \frac{1}{25}$ 	✓ expressing 125 as 5^3 ✓ raising both sides to reciprocal exponent of $-\frac{2}{3}$ ✓ answer (3)
2.2.2	$\frac{p^x \cdot p^{\frac{2}{x}}}{p^{5-x}} = 1$ $p^{x + \frac{2}{x} - 5 + x} = p^0$ $x + \frac{2}{x} - 5 + x = 0$ $2x^2 - 5x + 2 = 0$ $(2x - 1)(x - 2) = 0$ $x = \frac{1}{2} \quad \text{or} \quad x = 2$ OR $\frac{p^x \cdot p^{\frac{2}{x}}}{p^{5-x}} = 1$ $p^{x + \frac{2}{x}} = p^{5-x}$ $x + \frac{2}{x} = 5 - x$ $2x^2 - 5x + 2 = 0$ $(2x - 1)(x - 2) = 0$ $x = \frac{1}{2} \quad \text{or} \quad x = 2$	✓ laws of exponents ✓ equating bases ✓ standard form ✓ both answers (4) OR ✓ law of exponents ✓ equating bases ✓ standard form ✓ answers (4)


CAPS-GRADE 11
Marking Guidelines

2.2.3	$\frac{x-4}{\sqrt{x}-2} = 5a^0$ $\frac{(\sqrt{x}-2)(\sqrt{x}+2)}{\sqrt{x}-2} = 5(1)$ $\sqrt{x}+2 = 5$ $\sqrt{x} = 3$ $x = 9$ OR $\frac{x-4}{\sqrt{x}-2} = 5a^0$ $x-4 = 5(1)(\sqrt{x}-2)$ $x-4 = 5\sqrt{x}-10$ $(x+6)^2 = (5\sqrt{x})^2$ $x^2 + 12x + 36 = 25x$ $x^2 - 13x + 36 = 0$ $(x-9)(x-4) = 0$ $x = 9 \text{ or } x \neq 4$	$\checkmark (\sqrt{x}+2)(\sqrt{x}-2)$ $\checkmark a^0 = 1$ $\checkmark \sqrt{x} = 3$ $\checkmark \text{ answer} \quad (4)$ OR $\checkmark a^0 = 1$ $\checkmark \text{ isolating the root and squaring both sides}$ $\checkmark \text{ standard form}$ $\checkmark \text{ answer} \quad (4)$
[17]		

CAPS-GRADE 11
Marking Guidelines

QUESTION 3




Consider the following linear number pattern: $3x+8$; $x-2$; ... ; -141 ; -145 .

3.1	finite linear number pattern	✓ finite (1)
3.2	$3x+8-4=x-2$ $2x=-6$ $x=-3$	✓ $d = -4$ (from last 2 terms) ✓ substitution: $T_1 - 4 = T_2$ (2)
3.3	$-1 : -5 ; -9 : \dots$  $T_n = -4n + 3$	✓ first three terms ✓✓ answer (3)
3.4	$T_{15} = -4(15) + 3$ $T_{15} = -57$	✓ substitution of $n = 15$ ✓ answer (2)
3.5	$-145 = -4n + 3$ $4n = 148$ $n = 37$	✓ substitution of $T_n = -145$ ✓ answer (2)
[10]		

CAPS-GRADE 11
Marking Guidelines

QUESTION 4

Given a quadratic sequence: 3 ; 17 ; 35 ; 57 ; ...

4.1	next two terms: 83 and 113	✓ 83 ✓ 113 (2)
4.2	 $2a = 4 \therefore a = 2$ $3a + b = T_2 - T_1 \therefore 3(2) + b = 14 \therefore b = 8$ $a + b + c = T_1 \therefore 2 + 8 + c = 3 \therefore c = -7$ $\therefore T_n = 2n^2 + 8n - 7$ OR $2a = 4 \therefore a = 2$ $c = 3 + 4 - 14 = -7$ $\therefore T_n = 2n^2 + bn - 7$ $3 = 2(1)^2 + b - 7$ $b = 8$ $\therefore T_n = 2n^2 + 8n - 7$	✓ $2a = 4$ ✓ $3(2) + b = 14$ ✓ $2 + 8 + c = 3$ ✓ general term (4) OR ✓ $2a = 4$ ✓ $c = 3 + 4 - 14$ ✓ $3 = 2(1)^2 + b - 7$ ✓ general term (4)
4.3	$2n^2 + 8n - 7 > 826$ $2n^2 + 8n - 833 > 0$ CVs: $n = \frac{-(8) \pm \sqrt{(8)^2 - 4(2)(-833)}}{2(2)}$ $n = 18,51$ or $n \neq -22,51$ ($n \in \mathbb{N}$)   $n \geq 19$ $\therefore n = 20$ T_{20} is the second term greater than 826	✓ $T_n > 826$ ✓ critical values ✓ solving for n ✓ answer (4)


CAPS GRADE 11
Marking Guidelines

4.4	General term for the sequence of first differences: $T_n = 4n + 10$ $122 = 4n + 10$ $n = 28$ $\therefore T_{28}$ and quadratic seq. have a 1st difference of 122 OR $T_{n+1} - T_n = 122$ $2(n+1)^2 + 8(n+1) - 7 - (2n^2 + 8n - 7) = 122$ $2n^2 + 4n + 2 + 8n + 8 - 7 - 2n^2 - 8n + 7 = 122$ $4n = 112$ $n = 28$ $\therefore$ Between T_{28} and T_{29}	✓ $T_n = 4n + 10$ ✓ $122 = 4n + 10$ ✓ $n = 28$ ✓ T_{28} and T_{29} (4) OR ✓ $T_{n+1} - T_n = 122$ ✓ substitution ✓ $n = 28$ ✓ T_{28} and T_{29} (4)
4.5	$8p - 2 = -2(17)$ and $18q = -2(57)$ $8p - 2 = -34$ and $18q = -114$ $\therefore p = -4$ and $\therefore q = \frac{-19}{3}$ or $-6\frac{1}{3}$ or $-6,33$	✓ $p = -4$ ✓ $q = \frac{-19}{3}$ (2)
[16]		

CAPS-GRADE 11
Marking Guidelines

QUESTION 5

Given: $h(x) = \frac{-3}{x - \frac{3}{2}} - 2$



5.1.1	Equation of vertical asymptote: $x = \frac{3}{2}$ Equation of horizontal asymptote: $y = -2$	✓ $x = \frac{3}{2}$ ✓ $y = -2$ (2)
5.1.2	Coordinates of the y-intercept: $h(0) = \frac{-3}{2(0) - 3} - 2$ $y = -1$ $(0; -1)$ Answer only: full marks	✓ $x = 0$ ✓ $y = -1$ (2)
5.1.3	Coordinates of the x-intercept $0 = \frac{-3}{2x - 3} - 2$ $2 = \frac{-3}{2x - 3}$ $2(2x - 3) = -3$ $4x - 6 = -3$ $x = \frac{3}{4}$ $\left(\frac{3}{4}; 0\right)$	✓ $h(x) = 0$ ✓ simplification ✓ answer (3)
5.1.4		✓ both asymptotes ✓ both intercepts ✓ shape (3)

CAPS-GRADE 11
Marking Guidelines

5.1.5	Eqn. of axis of symmetry of h that has a negative gradient: $y = -x + c$ $-2 = -\left(\frac{3}{2}\right) + c$ $c = -\frac{1}{2}$ $y = -x - \frac{1}{2}$  Answer only: full marks	✓ substitution of gradient of -1 and point $\left(\frac{3}{2}; -2\right)$ ✓ answer (2)
5.1.6	Coordinates of points of $\cap$ of h and axis of symmetry: $\frac{-3}{x - \frac{3}{2}} - 2 = -x - \frac{1}{2}$ $\frac{-3}{x - \frac{3}{2}} = -x + \frac{3}{2}$ $\frac{-3}{x - \frac{3}{2}} \left(x - \frac{3}{2}\right) = \left(-x + \frac{3}{2}\right) \left(x - \frac{3}{2}\right)$ $-3 = \left(-x + \frac{3}{2}\right) \left(x - \frac{3}{2}\right)$ $-3 = -x^2 + 3x - \frac{9}{4}$ $4x^2 - 12x - 3 = 0$ $x = \frac{-(-12) \pm \sqrt{(-12)^2 - 4(4)(-3)}}{2(4)}$ $x = 3, 23 \quad \text{or} \quad x = -0, 23$ $y = -3, 73 \quad \text{or} \quad y = -0, 27$ $\therefore (-0, 23; -0, 27) \text{ and } (3, 23; -3, 73)$	✓ equating f and axis of symmetry ✓ standard form ✓ substitution into correct formula ✓ both x-values ✓ both y-values (5)
5.2.1	Equation of k: $q = 1$ $y = b^x + 1$ $\frac{3}{2} = b^1 + 1$ $b = \frac{1}{2}$ $k(x) = \left(\frac{1}{2}\right)^x + 1$ OR $k(x) = 2^{-x} + 1$	✓ value of q ✓ value of b ✓ equation of k (3)
5.2.2	Describe, in words, the transformation from k to $j(x) = 2^x$: Reflect k about the y-axis and translate 1 unit down (in any order)	✓ reflection about y-axis ✓ translation of 1 unit downwards (2)
		[22]

CAPS–GRADE 11
Marking Guidelines

QUESTION 6

$$f(x) = \frac{1}{2}(x-2)^2 - 8 \text{ and } g(x) = x - 6$$

6.1	$f(x) = \frac{1}{2}(x-2)^2 - 8$ $0 = \frac{1}{2}(x-2)^2 - 8$ $(x-2)^2 = 16$ $x-2 = \pm 4$ $x_A = 6 \text{ or } x_B = -2$ $AB = 6 - (-2)$ $AB = 8$ OR $f(x) = \frac{1}{2}(x-2)^2 - 8$ $0 = \frac{1}{2}(x-2)^2 - 8$ $0 = \frac{1}{2}x^2 - 2x - 6$ $x^2 - 4x - 12 = 0$ $(x-6)(x+2) = 0$ $x_A = 6 \text{ or } x_B = -2$ $AB = 6 - (-2)$ $AB = 8$	✓ $f(x) = 0$ ✓ square root on both sides with $\pm$ ✓ both values of x ✓ length of AB (4) OR ✓ $f(x) = 0$ ✓ standard form ✓ both values of x ✓ length of AB (4)
6.2	D(2 ; -8)	✓ answer (1)
6.3	Range of f: $y \geq -8$ OR $y \in [-8 ; \infty)$	✓ answer (1)
6.4	Average gradient between points A and C C(0 ; -6) $m = \frac{y_2 - y_1}{x_2 - x_1}$ $m = \frac{-6 - 0}{0 - (-2)}$ $m = -3$	✓ coordinates of C ✓ correct substitution into gradient formula ✓ answer (3)

CAPS–GRADE 11
Marking Guidelines

6.5	For 2 +ve roots of $\frac{1}{2}x^2 - 2x - 6 - k = 0$: $-8 < k < -6$ OR $x \in (-8; -6)$	✓✓ answer (2)
6.6.1	Values of x for which $f(x) \geq 0$: $x \leq -2$ or $x \geq 6$ OR $x \in (-\infty; -2] \cup [6; \infty)$	✓✓ answer (2)
6.6.2	Values of x for which $\frac{g(x)}{f(x)} > 0$: $x > -2$; $x \neq 6$ OR $x \in (-2; 6) \cup (6; \infty)$	✓✓ answer (2)
6.7	Maximum value of h if $h(x) = \sqrt{2}^{2-f(x)}$: $2-f(x)$ is a max. when $f(x)$ is a min. min. of $f(x) = -8$ $\therefore 2-f(x)$ has a max. value of 10 max. of $h(x) = (\sqrt{2})^{10}$ $= 2^5$ $= 32$	✓ min. value of $f(x) = -8$ ✓ max. value of exponent = 10 ✓ answer (3)
6.8	Values of k for which $y = -3x + k$ will not be a tan. to f: $\frac{1}{2}(x-2)^2 - 8 = -3x + k$ $\frac{1}{2}x^2 - 2x + 2 - 8 = -3x + k$ $x^2 + 2x - 12 - 2k = 0$ $a = 1$ $b = 2$ $c = -12 - 2k$ Line will be a tangent to f when $\Delta = 0$: $(2)^2 - 4(1)(-12 - 2k) = 0$ $4 + 48 + 8k = 0$ $k = -\frac{13}{2}$ $\therefore k \neq -6\frac{1}{2}$	✓ equating both equations ✓ standard form in terms of k ✓ substitution into $b^2 - 4ac$ ✓ value of k ✓ answer (5)

[23]


CAPS–GRADE 11
Marking Guidelines

QUESTION 7

7.1	$A = P(1-i)^n$ $A = R625\,000(1-0,1025)^6$ $A = R326\,653,08$	✓ substitution ✓ answer (2)
7.2	$1+i_{\text{eff}} = \left(1+\frac{i_{\text{nom}}}{m}\right)^m$ $i_{\text{eff}} = \left(1+\frac{0,08}{2}\right)^2 - 1$ $r_{\text{eff}} = \frac{51}{625} \times 100\%$ $r = 8,16\%$	✓ formula ✓ substitution ✓ answer (3)
7.3	$A = P(1+i)^n$ $1\,671\,650,74 = P\left(1+\frac{0,14}{4}\right)^{4 \times 4,5}$ $P = \frac{1\,671\,650,74}{\left(\frac{207}{200}\right)^{18}}$ $P = R899\,951,80$	✓ $n = 4 \times 4,5 = 18$ ✓ substitution into correct formula ✓ answer (3)
7.4.1	$75\,000 = P_3\left(1+\frac{0,08}{2}\right)^{1 \times 2}$ $P_3 = R69\,341,72$	✓ $n = 2$ ✓ $i = \frac{0,08}{2}$ or $\frac{8\%}{2}$ ✓ answer (3)

CAPS-GRADE 11
Marking Guidelines

7.4.2



$$75\,000 = \left[x \left(1 + \frac{0,12}{12}\right)^{36} - 15\,000 \left(1 + \frac{0,12}{12}\right)^{12} \right] \left(1 + \frac{0,08}{2}\right)^2$$

$$x = \frac{\frac{75\,000}{\left(1 + \frac{0,08}{2}\right)^2} + 15\,000 \left(1 + \frac{0,12}{12}\right)^{12}}{\left(1 + \frac{0,12}{12}\right)^{36}}$$

$$x = R60\,278,15$$

OR

$$\begin{aligned} \text{Amount in account after R1 500 was withdrawn} &= \frac{69\,341,72}{\left(1 + \frac{0,12}{12}\right)^{12}} \\ &= R61\,537,25569 \end{aligned}$$

$$\begin{aligned} \text{Amt. in acc. before R1500 was withdrawn} &= 61\,537,25569 + 15\,000 \\ \text{i.e. value after 2 years} &= R76\,537,25569 \end{aligned}$$

$$x \left(1 + \frac{0,12}{12}\right)^{24} = 76\,537,25569$$

$$x = \frac{76\,537,25569}{\left(1 + \frac{0,12}{12}\right)^{24}}$$

$$x = R60\,278,15$$

$$\checkmark x \left(1 + \frac{0,12}{12}\right)^{36}$$

$$\checkmark -15\,000 \left(1 + \frac{0,12}{12}\right)^{12}$$

$$\checkmark \frac{75\,000}{\left(1 + \frac{0,08}{2}\right)^2}$$

$$\checkmark \text{ answer} \quad (4)$$

OR

Amount after withdrawal:

$$\checkmark \frac{69\,341,72}{\left(1 + \frac{0,12}{12}\right)^{12}}$$

Amount before withdrawal

$$\checkmark R61\,537,2521 + R15\,000$$

$$\checkmark \text{ equation}$$

$$\checkmark \text{ answer}$$

(4)

[15]

CAPS–GRADE 11
Marking Guidelines


QUESTION 8

8.1.1	$P(A \text{ or } B) = P(A) + P(B)$ $= 0,35 + 0,40$ $= 0,75$	✓ $0,35 + 0,40$ ✓ answer (2)
8.1.2	$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$ $= 0,35 + 0,40 - 0,35 \times 0,40$ $= 0,61$	✓ $P(A \text{ and } B) = 0,35 \times 0,40 = 0,14$ ✓ correct substitution into rule ✓ answer (3)
8.2.1	$k = 18$	✓ answer (1)
8.2.2	$P(\text{Boy who does not listen to Pop music}) = \frac{32 + 23}{150}$ $= \frac{11}{30} \approx 0,37$	✓✓ answer (2)
8.2.3	$P(\text{Girl and listens to Pop music}) = \frac{18}{150}$ $= 0,12$ $P(\text{Girl}) \times P(\text{listens to Pop music}) = \frac{80}{150} \times \frac{33}{150}$ $= 0,12$ $P(\text{Girl and listens to Pop}) = P(\text{Girl}) \times P(\text{listens to Pop})$ $\therefore \text{The events are independent.}$	✓ $\frac{18}{150}$ or $0,12$ ✓ $\frac{80}{150} \times \frac{33}{150}$ ✓ $0,12$ ✓ conclusion (4)
		[12]

CAPS-GRADE 11
Marking Guidelines

QUESTION 9

9.1



$$0,35 - x + x + 0,60 - x + 0,08 = 1$$

$$x = 0,03$$

$$P(\text{only NZ will reach the finals}) = 0,6 - 0,03 \\ = 0,57$$

OR

$$P(\text{NZ or SA}) = P(\text{NZ}) + P(\text{SA}) - P(\text{NZ and SA}) \\ 1 - 0,08 = 0,35 + 0,6 - P(\text{NZ and SA})$$

$$P(\text{NZ and SA}) = 0,03$$

$$P(\text{only NZ will reach the finals}) = 0,6 - 0,03 \\ = 0,57$$

$$\checkmark 0,35 - x$$

$$\checkmark 0,60 - x$$

$$\checkmark \text{sum of probabilities} = 1$$

$$\checkmark P(\text{only NZ}) = 0,57 \quad (4)$$

OR


$$\checkmark P(\text{NZ or SA}) = 0,92$$

$$\checkmark \text{correct substitution into rule}$$

$$\checkmark P(\text{NZ and SA}) = 0,03$$

$$\checkmark P(\text{only NZ}) = 0,57 \quad (4)$$

9.2



$$\left(\frac{x}{x+8}\right)\left(\frac{8}{x+7}\right) + \left(\frac{8}{x+8}\right)\left(\frac{x}{x+7}\right) = \frac{48}{95}$$

$$\frac{8x + 8x}{(x+8)(x+7)} = \frac{48}{95}$$

$$\frac{x^2 + 15x + 56}{16x} = \frac{95}{48}$$

$$48x^2 + 720x + 2688 = 1520x$$

$$3x^2 - 50x + 168 = 0$$

$$x = \frac{-(-50) \pm \sqrt{(-50)^2 - 4(3)(168)}}{2(3)}$$

$$x = 12 \quad \text{or} \quad x = \frac{14}{3}$$

There are 12 apples in the basket

$$\checkmark P(A).P(B) = \left(\frac{x}{x+8}\right)\left(\frac{8}{x+7}\right)$$

$$\checkmark P(B).P(A) = \left(\frac{8}{x+8}\right)\left(\frac{x}{x+7}\right)$$

$$\checkmark P(A).P(B) + P(B).P(A) = \frac{48}{95}$$

$$\checkmark \text{standard form}$$

$$\checkmark \text{answer}$$

[09]

TOTAL: 150