



Basic Education

KwaZulu-Natal Department of Education
REPUBLIC OF SOUTH AFRICA

MATHEMATICS P1

PREPARATORY EXAMINATION

SEPTEMBER 2016

MEMORANDUM


**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

MARKS: 150

This memorandum consists of 12 pages.

QUESTION 1

1.1.1	$2x(3x-5) = 0$ $x = 0$ or $x = \frac{5}{3}$	AA✓✓ Answers	(2)
1.1.2	$x^2 - 3x = 7$ $x^2 - 3x - 7 = 0$ $x = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(1)(-7)}}{2(1)}$ $x = 4,54$ or $x = -1,54$	A✓ equation in std. form CA✓ substitution Penalise rounding off CACA✓✓ answers	(4)
1.1.3	$2x - 5\sqrt{x} = 3$ $2x - 3 = 5\sqrt{x}$ $4x^2 - 12x + 9 = 25x$ $4x^2 - 37x + 9 = 0$ $(4x-1)(x-9) = 0$ $x = \frac{1}{4}$ or $x = 9$ n/a OR $2x - 5x^{\frac{1}{2}} - 3 = 0$ $\left(2x^{\frac{1}{2}} + 1\right)\left(x^{\frac{1}{2}} - 3\right) = 0$ $x^{\frac{1}{2}} = \frac{-1}{2}$ or $x^{\frac{1}{2}} = 3$ No solution $x = 9$	A✓ isolating $5\sqrt{x}$ M✓ squaring both sides CA✓ standard form CA✓ factors CA✓ both answers CA✓ rejecting A✓ standard form CA✓ factors CA✓ $x^{\frac{1}{2}} = \frac{-1}{2}$ CA✓ $x^{\frac{1}{2}} = 3$ CA✓ answer CA✓ rejecting	(6)
1.1.4	$2^x(3x+1) < 0$ $2^x > 0$ for all $x \in R$ $\therefore 3x+1 < 0$ $x < -\frac{1}{3}$ OR [Graphically] 	A✓ $2^x > 0$ for all $x \in R$ A✓ $3x+1 < 0$ A✓ $x < -\frac{1}{3}$ A✓ sketch of $y = 2^x$ A✓ sketch of $y = 3x+1$ A✓ $x < -\frac{1}{3}$	(3)

1.2	$2^{99} \cdot 2^1 - 2^{99}$ $2^{99}(2 - 1)$ $= 2^{99}$	M✓ for taking 2^{99} as a common factor A✓ $(2 - 1)$ A✓ answer	(3)
1.3	$2x - y = 3 \rightarrow (1)$ $x^2 + 5xy + y^2 = 15 \rightarrow (2)$ From (1) : $y = 2x - 3$ Subst. (1) into (2) $x^2 + 5x(2x - 3) + (2x - 3)^2 = 15$ $x^2 + 10x^2 - 15x + 4x^2 - 12x + 9 - 15 = 0$ $15x^2 - 27x - 6 = 0$ $(5x + 1)(3x - 6) = 0$ $x = \frac{-1}{5} \text{ or } x = 2$ $y = \frac{-17}{5} \text{ or } y = 1$ OR $2x - y = 3 \rightarrow (1)$ $x^2 + 5xy + y^2 = 15 \rightarrow (2)$ From (1) : $x = \frac{y + 3}{2}$ Subst. (1) into (2) $\left(\frac{y + 3}{2}\right)^2 + 5y\left(\frac{y + 3}{2}\right) + y^2 = 15$ $\left(\frac{y^2 + 6y + 9}{4}\right) + \frac{5y^2 + 15y}{2} + y^2 = 15$ $y^2 + 6y + 9 + 10y^2 + 30y + 4y^2 = 60$ $15y^2 + 36y - 51 = 0$ $5y^2 + 12y - 17 = 0$ $(5y + 17)(y - 1) = 0$ $y = \frac{-17}{5} \text{ or } y = 1$ $x = \frac{-1}{5} \text{ or } x = 2$	A✓ y as subject CA✓ substitution $y = 2x - 3$ CA✓ std. form CA✓ factors CA✓ x values CA✓ y values A✓ x as subject CA✓ substitution $x = \frac{y + 3}{2}$ CA✓ std. form CA✓ factors CA✓ y values CA✓ x values	(6)
			[24]

[illegible]

QUESTION 3

3.1.1	6 ; 2 ; 10 ; 2 ; 14 ; 2 ; ... 18 ; 2	AA✓✓ answers	(2)
3.1.2	$a = 2$ and is a constant sequence $\therefore S_{50} = 50 \times 2$ $= 100$	A✓100	(1)
3.1.3	$a = 6; d = 4; n = 50$ $S_{50} = \frac{50}{2}[2(6) + 4(50 - 1)]$ $= 5200$ $S_{100} = 5200 + 100$ $= 5300$	A✓ for a, d and n values CA✓ for substitution into formula CA✓ 5200 CA✓ answer	(4)
3.2	$S_n = a + ar + ar^2 + \dots + ar^{n-1} \rightarrow (1)$ $rS_n = ar + ar^2 + \dots + ar^n \rightarrow (2)$ (2) – (1): $rS_n - S_n = ar^n - a$ $S_n(r - 1) = a(r^n - 1)$ $\therefore S_n = \frac{a(r^n - 1)}{r - 1}$	A✓ for equation (1) A✓ for equation (2) A✓ subtraction on LHS and RHS A✓ factorising if $S_n = \frac{a(1 - r^n)}{1 - r}$ penalize 1 mark	(4)
			[11]

QUESTION 4

4.1	$a = 1 \quad r = \frac{2}{3}$ $S_{\infty} = \frac{a}{1 - r}$ $= \frac{1}{1 - \frac{2}{3}}$ $= 3$	A✓ a and r value CA✓ substitution of r into formula CA✓ answer	(3)
4.2	$S_n = 3n^2 - n$ $T_4 = S_4 - S_3$ $= [3(4)^2 - (4)] - [3(3)^2 - (3)]$ $= 44 - 24$ $= 20$ OR	M✓ $T_4 = S_4 - S_3$ AA✓✓ substitution A✓ answer	(4)

	$S_1 = T_1 = a = 2$ $S_2 = T_1 + T_2 = 3(2)^2 - 2 = 10$ $T_2 = 10 - 2 = 8$ $S_3 = T_1 + T_2 + T_3 = 3(3)^2 - 3 = 24$ $T_3 = 24 - 8 - 2 = 14$ $d = 6$ $T_4 = a + 3d = 2 + 3(6) = 20$	A✓ a value A✓ second term value CA✓ d value CA✓ answer	(4)
			[7]

QUESTION 5

5.1	$A = P(1 - i)^n$ $A = 255000(1 - 0,125)^7$ $= R \quad 100 \quad 137,46$	A✓ formula A✓ correct substitution CA✓ answer	(3)
5.2.1	$A = P(1 + i)^n$ $= 10 \quad 000 \left(1 + \frac{0,095}{12}\right)^5$ $= R10 \quad 402,15$	A✓ correct substitution into correct formula CA✓ answer	(2)
5.2.2	$P = \frac{x[1 - (1 + i)^{-n}]}{i}$ $10 \quad 402,15 = \frac{450 \left[1 - \left(1 + \frac{0,095}{12}\right)^{-n}\right]}{\frac{0,095}{12}}$ $\frac{10 \quad 402,15 \times \frac{0,095}{12}}{450} = 1 - \left(1 + \frac{0,095}{12}\right)^{-n}$ $\left(1 + \frac{0,095}{12}\right)^{-n} = 0,816999213$ $-n = \frac{\log 0,816999213}{\log \left(1 + \frac{0,095}{12}\right)}$ $n = 25,6315128$ $\therefore n = 26 \text{ payments}$	CA✓ correct substitution of P into formula A✓ $x = 450$ CA✓ n as subject M✓ using logs CA($n \in \mathbb{N}$)✓ answer	(5)

5.2.3	Balance on Loan after the 25th payment $= 10\,402,15 \left(1 + \frac{0,095}{12}\right)^{25} - \frac{450 \left[\left(1 + \frac{0,095}{12}\right)^{25} - 1 \right]}{\frac{0,095}{12}}$ $= R12668,90 - R12386,54$ $= R282,36$ OR $P = \frac{x[1 - (1+i)^{-n}]}{i}$ $= \frac{450 \left[1 - \left(1 + \frac{0,095}{12}\right)^{-0,6315128} \right]}{\frac{0,095}{12}}$ $= R282,36$	M✓ (A - F_v) formula A✓ n = 25 CA✓ $10\,402,15 \left(1 + \frac{0,095}{12}\right)^{25}$ CA✓ $\frac{450 \left[\left(1 + \frac{0,095}{12}\right)^{25} - 1 \right]}{\frac{0,095}{12}}$ CA✓ answer OR A✓ formula A✓ subst. 450 CA✓ subst $\frac{0,095}{12}$ CA✓ subst 0,6315128 CA✓ answer	(5)
			(5)
			[15]

QUESTION 6 (For learners who did not receive the errata and used P(-4;4) , should be marked accordingly)


6.1	$p = -3$ $q = -2$	A✓ p value A✓ q value	(2)
6.2	$h(x) = \frac{a}{x+p} + q$. P(4;-4) is a point on h $-4 = \frac{a}{4-3} - 2$ $\therefore a = -2$	CA✓ subst. p, q and point P CA(negative)✓ a value	(2)
6.3	$h(x) = \frac{-2}{x-3} - 2$ $h(0) = \frac{-2}{0-3} - 2$ $= -1\frac{1}{3}$ $\left(0; -1\frac{1}{3}\right)$	A✓ substituting x = 0 CA(negative)✓ answer	(2)
6.4	x = 1	CACA✓✓ answer	(2)

6.5	$y = -(x + p) + q$ $y = -(x - 3) - 2$ $y = -x + 1$ $\therefore c = 1$ OR $y = -x + c$ <i>Point of intersection of asymptotes</i> $(3 ; -2)$ $-2 = -3 + c$ $c = 1$	CA✓substitution of p and q values into equation of line of symmetry CA✓answer CA✓substitution of $(3 ; -2)$ into equation of line of symmetry CA✓answer	 (2) (2)
			[10]

QUESTION 7

7.1	$D(0 ; -10)$	A(must be in coordinate form)✓ answer	(1)
7.2	$x^2 - 3x - 10 = 0$ $(x + 2)(x - 5) = 0$ $x = -2$ or $x = 5$ $A(-2;0)$ $B(5;0)$	A✓ $x^2 - 3x - 10 = 0$ CA✓factors CACA(negative and positive)✓✓ each x - value $A(-2;0)$ $B(5;0)$	(4)
7.3	$a = 2$ and $q = -10$	CA(positive)✓ a - value A✓ q - value	(2)
7.4	$x = -\frac{b}{2a} = -\frac{(-3)}{2(1)} = \frac{3}{2}$ or $x = \frac{-2+5}{2} = \frac{3}{2}$ $y = \left(\frac{3}{2}\right)^2 - 3\left(\frac{3}{2}\right) - 10 = -\frac{49}{4} / -12,25 / -12\frac{1}{4}$ $C(1,5 ; -12,25)$	A✓ $x = -\frac{b}{2a} = -\frac{(-3)}{2(1)} = \frac{3}{2}$ or CA✓ $x = \frac{-2+5}{2} = \frac{3}{2}$ CA✓substitution CA✓minimum value	(3)
7.5	$(-1,5 ; -9,25)$	CA✓ x - value CA✓ y - value	(2)
7.6	$x \geq \frac{3}{2}$ OR $g'(x).h'(x) \geq 0$ $(2x - 3).2 \geq 0$ $x \geq \frac{3}{2}$	CACA✓✓answer CA✓product of derivatives CA(positive)✓answer penalize 1 mark for incorrect notation	 (2) (2)
			[14]

QUESTION 8

8.1	$y = \log_3 x$	AA✓✓ answer	(2)
8.2		A✓ Shape of p and p^{-1} A✓ y – intercept of p A✓ x – intercept of p^{-1} A✓ point on each graph	(4)
8.3	$\log_3 x = 3$ $x = 27$ $0 < x \leq 27$	M✓ setting up equation CA✓ $x = 27$ CACA✓✓ for end points and inequality ANSWER ONLY full marks	(4)
			[10]

QUESTION 9 [Penalise once for notational error in questions 9, 10 and 11]

9.1	$f(x) = x^3$ $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $= \lim_{h \rightarrow 0} \frac{(x+h)^3 - x^3}{h}$ $= \lim_{h \rightarrow 0} \frac{x^3 + 3x^2h + 3xh^2 + h^3 - x^3}{h}$ $= \lim_{h \rightarrow 0} \frac{h(3x^2 + 3xh + h^2)}{h}$ $= \lim_{h \rightarrow 0} (3x^2 + 3xh + h^2)$ $= 3x^2$	A✓ formula A✓ substitution into correct formula CA✓ simplifying CA✓ factors CA✓ answer	(5)
9.2.1	$f(x) = x^{\frac{1}{2}} - 4x^{-2}$ $f'(x) = \frac{1}{2}x^{-\frac{1}{2}} + 8x^{-3}$	AA✓✓ exponential form CACA✓✓ answer	(4)

9.2.2	$\left(\frac{1}{2}x^2 - 3\right)^2 = y$ $y = \frac{1}{4}x^4 - 3x^2 + 9$ $\frac{dy}{dx} = x^3 - 6x$ OR $\frac{dy}{dx} = 2\left(\frac{1}{2}x^2 - 3\right) \cdot x$ $= x^3 - 6x$	A✓squaring both sides A✓simplifying CACA✓✓answers OR AA✓✓derivative using chain rule CACA✓✓answers	(4) (4)
			[13]

QUESTION 10

10.1.1	$f(-3) = (-3)^2 - 8 = 1$	A✓ $y = 1$ (y – value)	(1)
10.1.2	$f'(x) = 2x$ $m = f'(-3) = 2(-3) = -6$	A✓ value of gradient of tangent	(1)
10.1.3	Equation of tangent : - $y = mx + c$ $1 = -6(-3) + c$ $c = -17$ $y = -6x - 17$ OR $y - 1 = -6(x + 3)$ $y = -6x - 18 + 1$ $y = -6x - 17$	CA✓ substitution CA✓ answer CA✓ substitution CA✓ answer	(2) (2)
10.2.1	$f(x) = x^3 - 3x - 2$ $f'(x) = 3x^2 - 3 = 0$ $3(x+1)(x-1) = 0$ $x = -1 \quad \text{or} \quad x = 1$ $y = 0 \quad \text{or} \quad y = -4$ $A(-1; 0) \quad B(1; -4)$	A✓ derivative <u>and</u> equating to 0 CA✓ factors CA✓ x - values CA✓ y - values	(4)

10.2.2	$PQ = g(x) - f(x)$ $= (2x - 2) - (x^3 - 3x - 2)$ $= -x^3 + 5x$ $PQ' = -3x^2 + 5 = 0$ $x^2 = \frac{5}{3}$ $x = \pm \sqrt{\frac{5}{3}} = 1,29$ For $x = 1,29$ $y = -(1,29)^3 + 5(1,29) = 4,3$ Maximum Length of $PQ = 4,3 \text{ units}$	A✓ expression for PQ A✓ $PQ' = 0$ CA✓ x – values CA✓ y value for $x = 1,29$	(4)
10.2.3	$k = -4$ or $k = 0$	$k = 0$ A✓ $k = -4$ CA✓ answers	(2)
10.2.4	$f''(x) = 6x > 0$ $x > 0$ OR $x = \frac{x_1 + x_2}{2} = \frac{(-1) + (1)}{2} = 0$ $x > 0$ OR $x = -\frac{b}{3a} = -\frac{0}{3(1)} = 0$ $x > 0$	A✓ $6x$ A✓ $6x > 0$ A✓ answer A✓ midpoint formula CA✓ $x = 0$ CA✓ answer A✓ formula A✓ $x = 0$ A✓ answer	(3) (3) (3)
			[17]

QUESTION 11 (Ignore units in the question)

11.1	$V = s'(t) = 2t + 15 \text{ m/s}$	A✓ answer	(1)
11.2	$V = s'(25) = 2(25) + 15 = 65 \text{ m/s}$	CA✓ answer	(1)
11.3	$V = s'(t) = 2t + 15 \text{ m/s}$ $A = s''(t) = 2 \text{ m/s}^2$	CACA✓✓ 2	(2)
11.4	$A = s''(5) = 2 \text{ m/s}^2$	CA✓ answer (answer only 1 mark)	(1)
11.5	$s(t) = t^2 + 15t = 250$ $t^2 + 15t - 250 = 0$ $(t + 25)(t - 10) = 0$ $t = 10 \text{ s}$ $V = s'(10) = 2(10) + 15 = 35 \text{ m/s}$	A✓ equating s to 250 CA✓ factors CA✓ $t = 10$ CA✓ answer	(4)
			[9]

QUESTION 12

12.1.1	150	A✓answer	(1)
12.1.2(a)	$P(\text{Male}) = \frac{70}{150} = \frac{7}{15}$	A✓answer	(1)
12.1.2(b)	$P(\text{Eating Chocolate}) = \frac{80}{150} = \frac{8}{15}$	A✓answer	(1)
12.1.3	$P(\text{Male eating chocolate}) = \frac{45}{150} = 0,3$ $P(\text{Male}) \times P(\text{Eating chocolate})$ $= \frac{70}{150} \times \frac{80}{150} = \frac{56}{225} = 0,249$ $P(\text{Male and Eating chocolate}) \neq$ $P(\text{Male}) \times P(\text{Eating chocolate})$ Events are not independent	A✓ $P(\text{male eating chocolate})$ CA✓ $P(\text{Male}) \times P(\text{eating chocolate})$ value CA✓conclusion	(3)
12.2.1	$6 \times 7 \times 7 \times 7 = 2058$	A✓ $6 \times 7 \times 7 \times 7$ A✓2058	(2)
12.2.2	$6 \times 6 \times 5 \times 4 = 720$	A✓ $6 \times 6 \times 5 \times 4$ A✓720	(2)
12.2.3	Four digit codes divisible by 5:- $6 \times 7 \times 7 \times 1 = 252$ Probability of a four digit code divisible by 5 $= \frac{252}{2058} = \frac{7}{49} = 0,1428 = 14,28 \%$	A✓ $6 \times 7 \times 7 \times 1$ A✓252 CA✓answer	(3)
			[13]

TOTAL MARKS: 150