

GAUTENG PROVINCE
EDUCATION
REPUBLIC OF SOUTH AFRICA

PREPARATORY EXAMINATION
2015
MEMORANDUM

SUBJECT: MATHEMATICS P2/ WISKUNDE V2 (10612)

GAUTENG DEPARTMENT OF EDUCATION
PREPARATORY EXAMINATION – 2015

MATHEMATICS/WISKUNDE
(Second Paper/Tweede Vraestel)

MEMORANDUM

NOTE:

- If a candidate answered a QUESTION TWICE, mark the FIRST attempt ONLY.
- Consistent accuracy applies in ALL aspects of the memorandum.
- Penalise for rounding **only** in QUESTION 3.6
- Assuming answers/values in order to solve a problem is NOT ACCEPTABLE.
- S/R refers to STATEMENT as well as REASON

NOTA:

- As 'n kandidaat 'n vraag TWEEKEER beantwoord, merk slegs die EERSTE poging.
- Volgehoue akkuraatheid word in ALLE aspekte van die nasienmemorandum toegepas.
- Penalisering vir afronding word **slegs** in vraag 3.6 toegepas.
- Aanvaarding van antwoorde/waardes om 'n probleem op te los, is ONAANVAARBAAR.
- S/R verwys na BEWERING sowel as die REDE.


QUESTION / VRAAG 1

1.1		✓ $(\bar{x}; \bar{y})$ (22,6 ; 44,5) ✓ line of best fit <i>beste paslyn/lyn van beste passing</i>
		(2)
1.2.1	25 years /25 jaar	✓ answer / antwoord (1)
1.2.2	The reaction time according to the line of best fit for age 25 should be 51,46. The reaction time of this patient is ± 65. <i>Die reaksietyd volgens die lyn van beste passing vir die ouderdom van 25 moet 51,46 wees. Die reaksietyd vir hierdie pasiënt is ± 65.</i>	✓ answer / antwoord (1) Any other valid explanation. <i>Enige ander geldige verduideliking</i>
1.3	A = -21,03 B = 2,90 $y = 2,90x - 21,03$	✓ A = -21,03 ✓ B = 2,90 ✓ $y = 2,90x - 21,03$
		(3)
1.4	$r = 0,95$ very strong correlation / <i>baie sterk korrelasie</i>	✓ $r = 0,95$ ✓ very strong / strong <i>baie sterk / sterk</i>
		(2)
1.5	The 30 year olds have a higher reaction time on average. The interquartile range of this group is bigger (IQR = 15) than that of the 15 year olds (IQR=5). The reaction time of the 30 year olds varies more. <i>Die 30-jariges het gemiddeld 'n hoër reaksietyd. Die interkwartielvariasiewydte van hierdie groep is groter (IQR=15). as dié van die 15-jariges (IQR=5).. Die reaksietyd van die 30-jariges varieer meer.</i>	✓ IOR ✓ for comments / <i>vir kommentaar</i>
		(2)
		[11]

QUESTION / VRAAG 2


2.1	<table border="1"> <thead> <tr> <th>Age in years <i>Ouderdom in jare</i></th><th>Number of people <i>Aantal mense</i></th><th>Cumulative frequency <i>Kumulatiewe frekwensie</i></th></tr> </thead> <tbody> <tr> <td>$0 \leq A < 10$</td><td>20</td><td>20</td></tr> <tr> <td>$10 \leq A < 20$</td><td>130</td><td>150</td></tr> <tr> <td>$20 \leq A < 30$</td><td>152</td><td>302</td></tr> <tr> <td>$30 \leq A < 40$</td><td>92</td><td>394</td></tr> <tr> <td>$40 \leq A < 60$</td><td>86</td><td>480</td></tr> <tr> <td>$60 \leq A < 80$</td><td>18</td><td>498</td></tr> <tr> <td>$80 \leq A < 100$</td><td>2</td><td>500</td></tr> </tbody> </table>	Age in years <i>Ouderdom in jare</i>	Number of people <i>Aantal mense</i>	Cumulative frequency <i>Kumulatiewe frekwensie</i>	$0 \leq A < 10$	20	20	$10 \leq A < 20$	130	150	$20 \leq A < 30$	152	302	$30 \leq A < 40$	92	394	$40 \leq A < 60$	86	480	$60 \leq A < 80$	18	498	$80 \leq A < 100$	2	500	✓ all answers/ <i>alle antwoorde</i> (1)
Age in years <i>Ouderdom in jare</i>	Number of people <i>Aantal mense</i>	Cumulative frequency <i>Kumulatiewe frekwensie</i>																								
$0 \leq A < 10$	20	20																								
$10 \leq A < 20$	130	150																								
$20 \leq A < 30$	152	302																								
$30 \leq A < 40$	92	394																								
$40 \leq A < 60$	86	480																								
$60 \leq A < 80$	18	498																								
$80 \leq A < 100$	2	500																								
2.2		✓ plotting of points <i>plot van punte</i> ✓ joining of points <i>verbind van punte</i> ✓ grounding point at (0 ; 0) <i>grond punt by (0 ; 0)</i> (3)																								
2.3.1	26 years old (accept 25 – 27) <i>26 jaar oud (aanvaar 25-27)</i>	✓ answer <i>antwoord</i> (1)																								
2.3.2	$\frac{90}{500} \times 100$ = 18% The percentage of people 16 and above 18 is 82% <i>Die persentasie mense 16 jaar en ouer is 82%</i>	✓ 90 ✓ 18% ✓ 82% (3) [8]																								

QUESTION / VRAAG 3



3.1	$M\left(\frac{-3-5}{2}; \frac{9+1}{2}\right)$ $M(-4; 5)$	$\checkmark x = -4$ $\checkmark y = 5$	(2)
3.2	$AM^2 = (-4+3)^2 + (5-9)^2$ $= 1 + 16$ $= 17$ $\therefore r = \sqrt{17}$ OR/OF $AD^2 = (-5+3)^2 + (1-9)^2$ $= 4 + 64$ $= 68$ $\therefore AD = \sqrt{68}$ $\therefore \text{radius} = \frac{\sqrt{68}}{2}$ $= \sqrt{17}$	$DM^2 = (-5+4)^2 + (1-5)^2$ $= 1 + 16$ $= 17$ $\checkmark$ correct substitution into distance formula <i>korrekte vervanging in afstand formule</i> $\checkmark r = \sqrt{17}$ $\checkmark$ correct substitution into distance formula <i>korrekte vervanging in afstand formule</i> $\checkmark r = \sqrt{17}$	(2)
3.3	Yes, the circle will pass through point C <i>Ja, die sirkel gaan deur punt C</i> $\hat{B} = \hat{C} = 90^\circ$, AD is the diameter line subtends equal $\angle^s$. $\hat{B} = \hat{C} = 90^\circ$, AD die middellyn lynstuk onderspan gelyke $\angle^e$	$\checkmark$ yes / ja $\checkmark$ reason / rede	(2)

3.4	B(0 ; y) $m_{AB} \times m_{BD} = -1$ $\left(\frac{9-y}{-3-0} \right) \left(\frac{1-y}{-5-0} \right) = -1$ $(9-y)(1-y) = -15$ $9 - 10y + y^2 = -15$ $y^2 - 10y + 24 = 0$ $(y-6)(y-4) = 0$ $\therefore y = 6 \text{ or } y = 4 \quad \therefore B(0;6)$ OR/OF $AB^2 + BD^2 = AD^2$ $(9-y)^2 + (-3-0)^2 + (-5-0)^2 + (1-y)^2 = (-5+3)^2 + (1-9)^2$ $81 - 18y + y^2 + 9 + 25 + 1 - 2y + y^2 = 4 + 64$ $2y^2 - 20y + 48 = 0$ $y^2 - 10y + 24 = 0$ $(y-6)(y-4) = 0$ $y = 6 \text{ or } y = 4$ $\therefore B(0;6)$	✓ $m_{AB} \times m_{BD} = -1$ ✓ $\left(\frac{9-y}{-3-0} \right) / \left(\frac{1-y}{-5-0} \right)$ ✓ standard form standaardvorm ✓ factors / faktore ✓ B(0 ; 6) (5) ✓ $(9-y)^2 + (-3-0)^2 + (-5-0)^2 + (1-y)^2$ ✓ $(-5+3)^2 + (1-9)^2$ ✓ standard form/ Standaardvorm ✓ factors / faktore ✓ B(0 ; 6) (5)
3.5	$m_{AB} = \frac{9-6}{-3-0}$ $= -1$ $m_{\parallel} = -1$ $y - y_1 = m(x - x_1) \quad \text{OR/OF} \quad y = -x + c$ $y - 1 = -1(x + 5) \quad -5 = -1 + c$ $y - 1 = -x - 5 \quad c = -4$ $y = -x - 4 \quad y = -x - 4$	✓ $m_{\parallel} = -1$ ✓ substitution of (-1 ; 5) vervang van (-1 ; 5) ✓ $y = -x - 4$ (3)



3.6	$m_{AD} = \frac{1-9}{-5+3} = 4$ $\tan \theta = 4$ $\theta = 76^\circ$ $m_{DB} = \frac{1-6}{-5-0} = 1$ $\tan \alpha = 1$ $\alpha = 45^\circ$ $\hat{BDA} = 76^\circ - 45^\circ = 31^\circ$ ext $\angle$ of Δ OR/OR $AB^2 = (-3-0)^2 + (9-6)^2$ $AB = \sqrt{18}$ $BD^2 = (-5-0)^2 + (1-6)^2$ $BD = \sqrt{50}$ $\tan \hat{BDA} = \frac{\sqrt{18}}{\sqrt{50}} = 0,6$ $\hat{BDA} = 31^\circ$ 	$\checkmark m_{AD} = \frac{1-9}{-5+3} = 4$ $\checkmark \tan \theta = 4$ $\checkmark 76^\circ$ $\checkmark \tan \alpha = 1$ $\checkmark 45^\circ$ $\checkmark 31^\circ \text{ (-1 if NOT rounded)}$ $31^\circ \text{ (-1 as NIE afgerond)}$ $\checkmark (-3-0)^2 + (9-6)^2$ $\checkmark AB = \sqrt{18}$ $\checkmark (-5-0)^2 + (1-6)^2$ $\checkmark BD = \sqrt{50}$ $\checkmark \text{using a correct trig ratio}$ $\text{gebruik korrekte trig verh}$ $\checkmark 31^\circ \text{ (-1 if NOT rounded)}$ $31^\circ \text{ (-1 as NIE afgerond)}$ (6) [20]
-----	--	--

QUESTION / VRAAG 4

4.1.1	$x^2 + y^2 - 2x + 6y = 0$ $(x-1)^2 - 1 + (y+3)^2 - 9 = 0$ $(x-1)^2 + (y+3)^2 = 1+9$ $(x-1)^2 + (y+3)^2 = 10$ $\therefore \text{centre : } (1; -3)$ $\therefore \text{radius : } \sqrt{10}$	$\checkmark (x-1)^2 - 1$ $\checkmark (y+3)^2 - 9$ $\checkmark (1; -3)$ $\checkmark \sqrt{10}$ (4)
4.1.2	$m_{\text{radius}} = \frac{-3+4}{1+2} = \frac{1}{3}$ $\therefore m_r \times m_t = -1$ $\therefore m_t = -3$ equation of tangent / vergelyking van raaklyn $y+4 = -3(x+2) \quad \text{OR} \quad y = -3x + c$ $y+4 = -3x-6 \quad -4 = -3(-2) + c$ $y = -3x-10 \quad \therefore c = -10$ $y = -3x-10$	$\checkmark m_r = \frac{1}{3}$ $\checkmark m_t = -3$ $\checkmark \text{substitution of } (-2; -4)$ $\text{vervanging van } (-2; -4)$ $\checkmark y = -3x-10$ (4)

4.2.1	$RA = RB$ $RA^2 = RB^2$ $(p-2)^2 + (q-3)^2 = (p+1)^2 + (q-6)^2$ $p^2 - 4p + 4 + q^2 - 6q + 9 = p^2 + 2p + 1 + q^2 - 12q + 36$ $-6p + 6q = 24$ $\therefore p - q = -4$	$\checkmark RA^2 = RB^2$ $\checkmark (p-2)^2 + (q-3)^2$ $\checkmark (p+1)^2 + (q-6)^2$ $\checkmark -6p + 6q = 24$ (4)
4.2.2	$R(p; q)$ lies on the line / lê opdielyn $2x + 5y + 1 = 0$ $2p + 5q = -1 \dots\dots(1)$ $p - q = -4$ $p = q - 4 \dots\dots(2)$ OR sub(1) into (2) / verv (1) in (2) $2p - 2q = -8 \dots\dots(2)$ $2p - 2q = -8 \dots\dots(2)$ $2(q - 4) + 5q = -1$ $2p + 5q = -1 \dots\dots(1)$ $2q - 8 + 5q = -1$ $(2) - (1) \quad 7q = 7$ $7q = 7$ $q = 1$ $q = 1$ sub q into equation (2) / verv q in vgl (2) $p = 1 - 4$ $p = -3$ $R(-3; 1)$ radius = RA OR/OF radius = RB $RA^2 = (-3-2)^2 + (1-3)^2$ $RB^2 = (-3+1)^2 + (1-6)^2$ $= 25 + 4$ $= 4 + 25$ $RA = \sqrt{29}$ $RB = \sqrt{29}$ $(x+3)^2 + (y-1)^2 = 29$	$\checkmark 2p + 5q = -1$ $\checkmark 2(q-4) + 5q = -1 /$ $2p - 2q = -8$ $\checkmark q = 1$ $\checkmark p = -3$ $\checkmark RA^2 / RB^2$ $\checkmark RA / RB = \sqrt{29}$ $\checkmark (x+3)^2 + (y-1)^2 = 29$ (7) [19]

QUESTION / VRAAG 5

5.1.1	$3y - 2x = 0$ $3y = 2x$ $y = \frac{2}{3}x$ $\tan \alpha = m$ $\tan \alpha = \frac{2}{3}$	$\checkmark y = \frac{2}{3}x$ $\checkmark \tan \alpha = m$	(2)
5.1.2	 $OP^2 = 2^2 + 3^2$ <i>Pyth</i> $OP = \sqrt{13}$ $\sin \alpha = \frac{2}{\sqrt{13}}$ or / of $\frac{2\sqrt{13}}{13}$ (with rational denominator) (met rasionale noemer)	$\checkmark OP = \sqrt{13}$ $\checkmark \sin \alpha = \frac{2}{\sqrt{13}}$	(2)
5.2.1	 $\widehat{QOP} = \beta - \alpha$	$\checkmark \beta - \alpha$	(1)
5.2.2	$\sin \widehat{QOP} = \sin(\beta - \alpha)$ $= \sin \beta \cos \alpha - \cos \beta \sin \alpha$ $= \left(\frac{3}{5} \times \frac{3}{\sqrt{13}} \right) - \left(\frac{-4}{5} \times \frac{2}{\sqrt{13}} \right)$ $= \frac{17}{5\sqrt{13}}$	$\checkmark \sin(\beta - \alpha)$ $= \sin \beta \cos \alpha - \cos \beta \sin \alpha$ $\checkmark \frac{3}{5} \times \frac{3}{\sqrt{13}}$ $\checkmark \frac{-4}{5} \times \frac{2}{\sqrt{13}}$ $\checkmark \frac{17}{5\sqrt{13}}$	(4) [9]

QUESTION / VRAAG 6


6.1	$\frac{\cos(40^\circ - x) \cdot \cos x - \sin(40^\circ - x) \cdot \sin x}{\sin 205^\circ \cdot \cos 25^\circ}$ $= \frac{\cos(40^\circ - x + x)}{-\sin 25^\circ \cdot \cos 25^\circ}$ $= \frac{\cos(40^\circ - x + x)}{-\frac{1}{2}(2 \sin 25^\circ \cdot \cos 25^\circ)}$ $= \frac{\cos 40^\circ}{-\frac{1}{2} \sin 50^\circ}$ $= \frac{\cos 40^\circ}{-\frac{1}{2} \cos 40^\circ} \quad \text{OR/OF} \quad \frac{\sin 50^\circ}{-\frac{1}{2} \sin 50^\circ}$ $= -2$	$\checkmark \cos(40^\circ - x + x)$ $\checkmark -\sin 25^\circ$ $\checkmark -\frac{1}{2} \sin 50^\circ$ $\checkmark \sin 50^\circ = \cos 40^\circ \text{ or/of}$ $\cos 40^\circ = \sin 50^\circ \text{ in numerator}$ in noemer $\checkmark -2$ (5)
6.2.1	$\text{LHS/LK} = \frac{\cos^2 x - \sin^2 x}{\cos x + \sin x}$ $= \frac{(\cos x - \sin x)(\cos x + \sin x)}{\cos x + \sin x}$ $= \cos x - \sin x$ $= \text{RHS/RK}$	$\checkmark \cos 2x = \cos^2 x - \sin^2 x$ $\checkmark \text{factorising /faktorisering}$ (2)
6.2.2	$\cos x \left(\frac{\cos 2x}{\cos x + \sin x} \right) = \frac{1}{2}$ $\cos x (\cos x - \sin x) = \frac{1}{2}$ $\cos^2 x - \cos x \sin x = \frac{1}{2}$ $2\cos^2 x - 2\cos x \sin x = 1$ $2\cos^2 x - 1 = 2\cos x \sin x$ $\cos 2x = \sin 2x$	$\checkmark \cos x - \sin x$ $\checkmark \cos^2 x - \cos x \sin x = \frac{1}{2}$ $\checkmark 2\cos^2 x - 1 = \cos 2x$ $\checkmark 2\cos x \sin x = \sin 2x$ (4)
6.2.3	$\cos x \left(\frac{\cos 2x}{\cos x + \sin x} \right) = \frac{1}{2}$ $\cos 2x = \sin 2x$ $\tan 2x = 1$ $2x = 45^\circ + k \cdot 180^\circ ; k \in \mathbb{Z}$ $x = 22,5^\circ + k \cdot 90^\circ ; k \in \mathbb{Z}$ OR/OF	$\checkmark \tan 2x = 1$ $\checkmark 2x = 45^\circ + k \cdot 180^\circ$ $\checkmark x = 22,5^\circ + k \cdot 90^\circ ; k \in \mathbb{Z}$ (3)

	$\cos 2x = \sin 2x$ $\cos 2x = \cos(90^\circ - 2x)$ $2x = 90^\circ - 2x + k.360^\circ \quad \text{or} \quad 2x = 2x - 90^\circ + k.360^\circ ;$ $k \in \mathbb{Z}$ $4x = 90^\circ + k.360^\circ ; k \in \mathbb{Z}$ $x = 22,5^\circ + k.90^\circ ; k \in \mathbb{Z}$	$\checkmark \cos 2x = \cos(90^\circ - 2x)$ $\checkmark 4x = 90^\circ + k.360^\circ$ $\checkmark x = 22,5^\circ + k.90^\circ$ (3)
6.3.1	Area $\Delta ABC = \frac{1}{2}mn \sin 4x$	$\checkmark \text{Area } \Delta ABC = \frac{1}{2}mn \sin 4x$ (2)
6.3.2	Max area if $\sin 4x = 1$ $4x = 90^\circ$ $x = 22,5^\circ$	$\checkmark \sin 4x = 1$ $\checkmark x = 22,5^\circ$ (2)
6.3.3	Right-angled triangle/Reghoekige driehoek	$\checkmark$ answer
[18]		


QUESTION / VRAAG 7


7.1	$\sin(60^\circ - \theta) = 1$ $60^\circ - \theta = 90^\circ$ $\theta = -30^\circ$	$\checkmark \theta = -30^\circ$ (1)
7.2	period / periode = $\frac{360^\circ}{2}$ $= 180^\circ$	$\checkmark 180^\circ$ (1)
7.3	$-2 \leq y \leq 0$ of / or $[-2;0]$	$\checkmark -2 \leq y \leq 0$ (1)
7.4	$45^\circ < x < 135^\circ$ or/or $150^\circ < x \leq 180^\circ$	$\checkmark 45^\circ$ and 135° $\checkmark 150^\circ$ and 180° $\checkmark$ ALL inequalities correct <i>ALLE ongelykhede korrek</i> Subtract 1 mark for extra intervals Trek 1 punt af vir ekstra intervale (3)
7.5	$90^\circ < x < 150^\circ$	$\checkmark 90^\circ$ and/en 150° $\checkmark$ inequalities / ongelykhede (2)
[8]		

QUESTION / VRAAG 9




S/R	Statement and reason	S/R	Bewering en rede
9.1.1	(a) $AB \parallel CD$ corr. $\angle^s$ equal / ooreenk $\angle^c$ gelyk (b) line from centre $\perp$ to chord / Midpt. $\odot$; Midpt. koord (c) Radius	✓ reason / rede ✓ reason / rede ✓ reason / rede	(3)
9.1.2	Join OB / Verbind OB HB = 12 and/en GD = 5 (line from centre $\perp$ to chord / Midpt. $\odot$; Midpt. koord) In $\triangle OHB$ $(OH)^2 = (OB)^2 - (BH)^2 \quad \hat{H} = 90^\circ \text{ (Pyth)}$ $= (13)^2 - (12)^2$ $= 25$ $OH = 5$ In $\triangle OGD$ $(OG)^2 = (OD)^2 - (GD)^2$ $= (13)^2 - (5)^2$ $= 144$ $OG = 12$ $GH = OG - OH$ $= 12 - 5$ $= 7$	✓ S/R ✓ Pythagoras ✓ OH = 5 ✓ OG = 12 ✓ GH = 7	(5)

9.2.1	Tangents from same point / Tans from common pt <i>Raaklyne vanuit dies. punt</i> 	✓ reason /rede (1)
9.2.1	$\hat{B}_1 + \hat{B}_2 = 90^\circ$ (tan $\perp$ rad / raaklyn $\perp$ radius) $\hat{A}_1 + \hat{A}_2 = 90^\circ$ (tan $\perp$ rad / raaklyn $\perp$ radius) $\therefore$ AOBT is a cyclic quadrilateral/is 'n koordevierhoek (opposite $\angle$'s supplementary/oorstaande $\angle$'e suppl)	✓ S/R ✓ S/R ✓ R (3)
9.2.2	$OA = OB$ (radii) $\hat{T}_1 = \hat{T}_2$ (equal chords subtend equal angles/ <i>gelyke hoeke onderspan deur gelyke koorde</i>	✓ $OA = OB$ (radii) ✓ R (2)
9.2.3	$\hat{A}_2 + \hat{B}_2 = 110^\circ$ (ext $\angle$ of Δ / buite $\angle$ van Δ) But $\hat{A}_2 = \hat{B}_2$ ($\angle$'s opp equal sides / $\angle$'e teenoor gelyke sye) $\hat{A}_2 = 55^\circ$ or $\hat{B}_2 = 55^\circ$ $\hat{A}_2 = \hat{B}_2 = \hat{C} = 55^\circ$ (tan chord / $\angle$ tussen raaklyn en koord) OR/OF $\hat{O}_1 + \hat{O}_2 = 110^\circ$ (ext $\angle$ of cycl quad/buite $\angle$ van kvh) $\hat{O}_1 + \hat{O}_2 = 2\hat{C}$ ($\angle$ at centre = 2x$\angle$ at circ /midpts$\angle$ = 2xomtreks$\angle$) $\hat{C} = 55^\circ$	✓ S/R ✓ S/R ✓ $\hat{C} = 55^\circ$ ✓ S/R (4) ✓ S/R ✓ S ✓ R ✓ $\hat{C} = 55^\circ$ (4) [17]




QUESTION / VRAAG 10

10.1	Construction: Join OD and OB <i>Konstruksie: Verbind OD en OB</i> $\hat{O}_1 = 2\hat{A}$ ($\angle$ at centre = $2 \times \angle$ at circumference / Midpts $\angle = 2 \times$ Omtreks $\angle$) $\hat{O}_2 = 2\hat{C}$ ($\angle$ at centre = $2 \times \angle$ at circumference / Midpts $\angle = 2 \times$ Omtreks $\angle$) $\hat{O}_1 + \hat{O}_2 = 360^\circ$ ($\angle^s$ round a pt OR $\angle^s$ in a rev $\angle^e$ om 'n punt OF omwenteling $2\hat{A} + 2\hat{C} = 360^\circ$ $2(\hat{A} + \hat{C}) = 360^\circ$ $\hat{A} + \hat{C} = 180^\circ$ M	✓ Construction / Konstruksie ✓ S/R ✓ S/R ✓ S/R ✓ $2\hat{A} + 2\hat{C} = 360^\circ$ (5)
------	--	--



10.2.1	$\hat{A} = 180^\circ - y$ $\hat{B} = 180^\circ - x$ $\hat{C} = 180^\circ - z$	(opp $\angle^s$ of cyclic quad / teenoorst. $\angle^e$ van kvh)	✓ all 3 statements al 3 bewerings ✓ reason rede	(2)
10.2.2	$(180^\circ - y) + (180^\circ - x) + (180^\circ - z)$ $= 540^\circ - (x + y + z)$ $= 540^\circ - 180^\circ$ $= 360^\circ$	(sum of $\angle^s$ of $\Delta = 180^\circ$)	✓ substitution of angles vervanging van hoeke ✓ simplification vereenvoudiging ✓ 180° ✓ answer / antw	(4) [11]

QUESTION / VRAAG 11



11.1	$\hat{R}_2 = \hat{M}_2$ $\hat{R}_1 = \hat{T}$	(alt $\angle^s$ RE $\parallel$ TM / verw. $\angle^e$; RE $\parallel$ TM) corr $\angle^s$ RE $\parallel$ TM / ooreenk. $\angle^e$ RE $\parallel$ TM)	✓ (S) ✓ (R) ✓ (S) ✓ (R)	(4)
11.2	$\frac{EM}{EG} = \frac{RT}{RG}$ (line $\parallel$ one side of Δ OR prop theorem; RE $\parallel$ TM) (lyn $\parallel$ een sy van Δ) But / Maar RT = RM ($\angle^s$ opp equal sides ; $\hat{M}_2 = \hat{T}$ / $\angle^e$ teenoor gelyke sye ; $\hat{M}_2 = \hat{T}$) $\frac{EM}{EG} = \frac{RM}{RG}$		✓ (S) ✓ (R) ✓ (S) ✓ (R)	(4)

11.3	In $\triangle GYE$ en $\triangle GER$ $\hat{G} = \hat{G}$ (common /gemeenskaplik) $\hat{E}_1 = \hat{R}_1$ (given /gegee) $\hat{Y}_1 = \hat{E}_1 + \hat{E}_2$ (3rd $\angle$ of $\triangle$ / 3e $\angle$ van $\triangle$) $\triangle GYE \parallel \triangle GER$ ($\angle, \angle, \angle$)	✓ $\hat{G} = \hat{G}$ (S/R) ✓ $\hat{E}_1 = \hat{R}_1$ (S/R) ✓ $\hat{Y}_1 = \hat{E}_1 + \hat{E}_2$ (S/R) ✓ equiangular $\triangle$ /gelykhoekige $\triangle$ or/of $\angle, \angle, \angle$ (R) (4)
11.4	$\frac{EG}{EY} = \frac{RG}{RE} = \frac{YG}{EG}$ (equiang $\triangle$'s / $\triangle GYE \parallel \triangle GER$) $\therefore \frac{EG}{EY} = \frac{RG}{RE}$	✓ (R) equiang $\triangle$'s / $\triangle GYE \parallel \triangle GER$ gelyk $\triangle$'s $\triangle GYE \parallel \triangle GER$ (1)
11.5.1	$(RG)^2 = (GM)^2 - (RM)^2$ $\hat{GRM} = 90^\circ$ Pythag. $= (10)^2 - (6)^2$ $= 64$ $\therefore RG = 8$	✓ substitute into Pyth vervang in Pyth ✓ $RG = 8$ (2)
11.5.2	$\frac{GE}{GM} = \frac{GR}{GT}$ (line $\parallel$ one side of $\triangle$ OR prop theorem; $RE \parallel TM$) (lyn $\parallel$ een sy van $\triangle$) $\frac{GE}{10} = \frac{8}{14}$ $GE = 5\frac{5}{7}$ or/of $\frac{40}{7}$ or/of 5,71 units/eenhede	✓ $\frac{GE}{GM} = \frac{GR}{GT}$ /reason /rede ✓ $GT = 14$ ✓ substitution of values vervanging van waardes ✓ answer /antwoord (4) [19]

TOTAL/TOTAAL: 150

